

**Amendments to the Specification**

Following the abstract, please insert the uploaded Sequence Listing filed herewith.

Please replace paragraph [0065] of the published application with the following amended paragraph:

**[0065]** FIG. 11 shows EV1 binding inhibited in the presence of anti- $\alpha_2\beta_1$ . Binding of [ $^{35}\text{S}$ ]-methionine labeled EV1 to ovarian cancer cell lines in the presence and absence of either anti- $\alpha_2\beta_1$  or anti-DAF MAbs. Levels of [ $^{35}\text{S}$ ]-methionine labeled virus bound was determined by liquid scintillation counting on a 1450 ~~Microbeta~~ MICROBETA® TRILUX (Wallac, Finland).

Please replace paragraph [0070] of the published application with the following amended paragraph:

**[0070]** FIG. 16 shows binding of [ $^{35}\text{S}$ ]-methionine labeled EV1 to SkMel28 melanoma cells in the presence and absence of either anti- $\alpha_2\beta_1$  or anti-DAF MAbs. Levels of [ $^{35}\text{S}$ ]-methionine labeled virus bound was determined by liquid scintillation counting on a 1450 ~~Microbeta~~ MICROBETA® TRILUX (Wallac, Finland).  $\alpha_2\beta_1$  blockade resulted in significant inhibition of EV1 binding. Results are expressed as the mean of triplicate samples  $\pm$  standard error.

Please replace paragraph [0103] of the published application with the following amended paragraph:

**[0103]** Enteroviral receptor surface expression on cancer cells was analysed by flow cytometry. Dispersed cells ( $1 \times 10^6$ ) were incubated for 20 minutes on ice with the appropriate MAb (5  $\mu\text{g/ml}$  diluted in PBS) for 20 minutes. Cells were washed with PBS and pelleted by centrifugation before resuspension in 100  $\mu\text{l}$  of 1:50 dilution of R-phycoerythrin-conjugated  $\text{F(ab')}_2$  fragment of goat anti-mouse immunoglobulin (Dako, A/S, Denmark). Cells were again incubated on ice for 20 minutes, washed, pelleted and resuspended in PBS prior to flow

cytometric analysis. Cell surface receptor expression was analysed using a FACStar™ Analyser (Becton Dickinson, Sydney, Australia).

Please replace paragraph [0110] of the published application with the following amended paragraph:

**[0110]** Six-well tissue culture plates containing confluent monolayers of DOV13 cells were inoculated with 500 µl EV1 (multiplicity of infection [moi]= $10^5$  TCID<sub>50</sub>/ml) for 1 hour at 37°C. Unbound virus was removed by washing three times with methionine/cysteine free DMEM (ICN Biomedical, Ohio, USA) and cell monolayers were incubated in 1.3 ml of this media for a further 2 hours at 37° C. before addition of 300 µCi of [<sup>35</sup>S]-methionine translabel (ICN Biomedical, Ohio, USA). Infected monolayers were incubated overnight at 37° C. in a 5% CO<sub>2</sub> environment. Following three freeze/thaw cycles viral lysates were purified in a 5-30% sucrose gradient by velocity centrifugation for 95 minutes at 36,000 rpm in a Beckman XL-90 ultracentrifuge (SW41ti Rotor). Fractions were collected from the bottom of each tube and monitored by liquid scintillation counting (Wallac 1450' Microbeta MICROBETA® TRILUX, Finland) to locate 160S viral peak fraction used in viral binding assays.

Please replace paragraph [0113] of the published application with the following amended paragraph:

Approximately  $1 \times 10^6$  cells resuspended in 800 µl of RPMI containing 1% bovine serum albumin (BSA) were incubated in the presence of 20 µg/ml of MAb (anti- $\alpha_2\beta_1$  or anti-DAF diluted in PBS) for 1 hour at 4° C[[.]] followed by the addition of 300 µl ( $1 \times 10^6$ ) of [<sup>35</sup>S]-methionine labeled 160S EV1. After incubation at 4° C[[.]] for 2 hours cells were washed four times with serum free media and cell pellets dissolved in 200 µl 0.2M NaOH-1% SDS before the level of [<sup>35</sup>S]-methionine labeled virus bound was determined by liquid scintillation counting from triplicate samples. (Wallac 1450 ~~Microbeta~~ MICROBETA® TRILUX, Finland). Results were expressed as means±SE.

Please replace paragraph [0115] of the published application with the following amended paragraph:

[0115] [<sup>35</sup>S]-methionine labeled viral fractions were analysed by polyacrylamide gel electrophoresis (PAGE) and visualised by autoradiography. [<sup>35</sup>S]-methionine labeled 160S EV1 fractions were incubated with sample reducing buffer (250 mM TRIS, 0.2 g w/v SDS, 20% v/v glycerol, 10% v/v 2-mercaptoethanol and 0.01% w/v bromophenol blue, pH 6.8) for 10 minutes at 95° C[.] ~~deraturing~~ denaturing the virion. Denatured 160S vial peak fractions were then separated on a 15% Tris-HCl precast gel (~~BIORAD Ready-Gel~~ Bio-Rad READY GEL®, CA, USA) in conjunction with a Benchmark prestained midrange protein ladder (GIBCO, USA) at 180 V for 45 minutes. Visualisation of the major structural proteins and analysis of viral purity was by autoradiography on Hyperfilm MP (Amersham International, England) after 96 hour exposure.

Please replace paragraph [0117] of the published application with the following amended paragraph:

[0117] Cell suspensions of human peripheral blood lymphocytes, OVHS-1 and DOV-13 cells were challenged with EV1 (moi=1.0 ~~TCID<sub>50</sub>~~ TCID<sub>50</sub>/cell) and incubated for 24 h at 37° C. Levels of cell cytolysis were calculated as a function of release of LDH (a stable cytosolic enzyme that is released upon cell lysis), assessed by using a ~~Cyto-Tox~~ CYTOTOX 96® kit (Promega Corp. Maddison, Wis. USA) as per the manufactures instructions.

Please replace paragraph [0119] of the published application with the following amended paragraph:

[0119] DOV-13 cells were seeded in a 24-well plate at 500 or 5000 cells per well in 1 ml of RPMI 1640 containing 5% FCS onto a semi-solid 0.5% agarose layer. Cells were incubated for 48 h at ~~37°C~~ 37°C in a 5% ~~CO<sub>2</sub>~~ CO<sub>2</sub> atmosphere to allow spheroids to form, before the addition of EV1 (10<sup>5</sup> TCID<sub>50</sub>).

Please replace paragraph [0123] of the published application with the following amended paragraph:

[0123] Sera from infected mice were analyzed for viremia using real-time quantitative RT-PCR. Briefly, viral RNA was extracted from 10 µl of serum using a QIAamp® Viral RNA mini kit (Qiagen, Clifton Hill, Victoria, Australia) and eluted in a final volume of 40 µl according to manufacturer's instructions. Primers and probe for determination of EV1 viral RNA levels were designed using the ~~Primer Express~~ PRIMER EXPRESS™ 1.5 software (Applied Biosystems, Foster City, Calif., USA) and were based on the previously published EV1 sequence (Genbank accession number AF029859); forward ~~primer~~ primer (5'-CAAGACAGGGACCAAAGAGGAT-3') (SEQ ID NO: 1), reverse primer (5'-CCACTCGCCTGGTTGTAATCA-3') (SEQ ID NO: 2) and 6-FAM-labeled MGB-probe (5'-CCAATAGCTTCAACAATT-3') (SEQ ID NO: 3). One-step RT-PCR was performed using ~~Platinum~~ PLATINUM® Quantitative RT-PCR ~~ThermoScript~~ THERMOSCRIPT™ One-Step System on an ABI 7000 sequence detector. For generation of the standard curve, 10-fold dilutions of EV1 viral stock ( $1 \times 10^6$  TCID<sub>50</sub>/ml) was amplified with optimized concentration of primers and probe. In a volume of 25 µl, the reaction mixture comprised ~~[.];~~ 1x ~~ThermoScript~~ THERMOSCRIPT™ reaction mix, 500 nM forward, 900 nM reverse primer, 250 nM probe, 500 nM ROX, 0.5 µl ~~ThermoScript~~ THERMOSCRIPT™ Plus/Platinum Taq Mix and 5 µl extracted RNA. Thermal cycling conditions were subjected to 30 min at 60° C ~~[.];~~, followed by 5 min at 95° C ~~[.];~~ and then 40 cycles of 15 s at 95° C ~~[.];~~ and 1 min at 60° C.